PATIENT INFORMATION
HOVILPRI (hoe-vil-pree)
(pritelivir)
tablets, for oral use

What is HOVILPRI?

HOVILPRI is a prescription medicine used in adults with weakened immune systems (immunocompromised) to treat lesions caused by herpes simplex virus (HSV) infection that have not improved with the medicines acyclovir, valacyclovir, or famciclovir.

HOVILPRI does not cure HSV infection.

It is not known if HOVILPRI is safe and effective in children less than 18 years of age.

Before taking HOVILPRI, tell your healthcare provider about all of your medical conditions, including if you:

- are pregnant or plan to become pregnant. It is not known if HOVILPRI will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if HOVILPRI passes into your breast milk. Talk with your healthcare provider about the best way to feed your baby if you take HOVILPRI.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. HOVILPRI may affect the way other medicines work, and other medicines may affect the way HOVILPRI works.

Especially tell your healthcare provider if you take medicines used to lower stomach acid, including:

- proton pump inhibitors (PPIs) (such as esomeprazole, rabeprazole, omeprazole, lansoprazole, or pantoprazole).
- H2-receptor blockers (such as famotidine, ranitidine).
- antacids (medicines such as Tums, Mylanta, Maalox).

See **"How should I take HOVILPRI?"** and **"What should I avoid while taking HOVILPRI?"** if you take any of these medicines.

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

How should I take HOVILPRI?

- Take HOVILPRI exactly as your healthcare provider tells you to take it.
- The usual dose of HOVILPRI is:
  - On Day 1 (first day of treatment): take 400 mg by mouth (four 100 mg tablets) at one time.
  - After Day 1: take 100 mg by mouth (one 100 mg tablet) 1 time a day.
- Take HOVILPRI with or without food.
- Take HOVILPRI as prescribed by your healthcare provider until your HSV lesions have healed.
- **If you take medicines that reduce stomach acid:**
  - If you take an H2-receptor blocker (such as famotidine, ranitidine), take HOVILPRI at least 2 hours before or 12 hours after taking the H2-receptor blocker.
  - If you take an antacid (such as Tums, Mylanta, Maalox), take HOVILPRI 2 hours before or 2 hours after taking the antacid.
- If you miss a dose of HOVILPRI, take it as soon as you remember. If you don't remember until the next day's scheduled dose, **take 2 tablets** at that time.
- If you take too much HOVILPRI, call your healthcare provider, or the Poison Help line (1-800-222-1222), or go to the nearest hospital emergency room right away.

What should I avoid while taking HOVILPRI?

Avoid taking the following proton pump inhibitor medicines during treatment with HOVILPRI:

- esomeprazole or rabeprazole
- omeprazole or lansoprazole (daily dose above 20 mg)
- pantoprazole (daily dose above 40 mg)

Talk to your healthcare provider before taking any of these medicines.

What are the possible side effects of HOVILPRI?

The most common side effect of HOVILPRI is headache.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store HOVILPRI?

- Store HOVILPRI tablets between 68ºF to 77°F (20ºC to 25º).

Keep HOVILPRI and all medicines out of the reach of children.

General information about the safe and effective use of HOVILPRI.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use HOVILPRI for a condition for which it was not prescribed. Do not give HOVILPRI to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about HOVILPRI that is written for health professionals.

What are the ingredients in HOVILPRI?

Active ingredient: pritelivir mesylate

Inactive ingredients: colloidal silica dioxide, croscarmellose sodium, magnesium stearate, mannitol, and microcrystalline cellulose. The film-coat contains hydroxy propyl methyl cellulose and polyethylene glycol, and calcium diphosphate.

Distributed by: Asahi Kasei Therapeutics, Inc. Cary, North Carolina 27518

HOVILPRI® is a registered trademark of Asahi Kasei Therapeutics, Inc. or its subsidiary

For more information, go to www.HOVILPRI.com or call 1-844-835-6947

This Patient Information has been approved by the U.S. Food and Drug Administration. Issued: 10/2026